

when males of two biotypes were present with a female of a given biotype in the same arena, the B male and female responded by increasing their frequency of courtships, leading to more copulation events, whereas the ZHJ1 male and female did not do so. Moreover, although courtships between the two biotypes occurred, copulation never resulted; this confirmed that both B and ZHJ1 share incompletely isolated mate recognition systems. Further, B males not only courted females of either biotype more frequently than did ZHJ1 males, they also interfered more frequently with the courtships initiated by ZHJ1 males than did ZHJ1 males with courtships initiated by B males (tables S2 and S3). The mating behavior and interactions between B and AN differed in ways similar to what we found for B and ZHJ1, although details varied between the two combinations (J8) (tables S4 and S5).

These results help to explain the underlying basis of the B biotype's capacity to invade and displace indigenous populations. The strong competitive ability of B results partly from its capacity to adjust sex ratio in favor of its population increase, and partly from its capacity to interfere with the mating of indigenous individuals. When the proportion of males is increased, B adults respond by increasing the frequency of copulation and consequently increasing the proportion of female progeny. Critical to this is that B responds independently of whether the males are all B or a mix. This interaction is extraordinary because the indigenous males actually help to promote copulation among the invaders and consequently increase the invaders' competitive capacity. In contrast, the indigenous females do not respond to increased numbers of adult males. Moreover, copulation by indigenous individuals is partly blocked by B males that readily attempt to court with females of either biotype—a behavior not reciprocated by the indigenous males. These asymmetric mating interactions have obvious population-level implications because the increase in the proportion of B females and the concomitant decrease in the proportion of indigenous females results in an immediate higher population growth rate for B and a lower growth rate for the indigenous population. As the abundance of B increases relative to the indigenous individuals, the increased allocation of eggs to female progeny and the active interference of mating of indigenous males by B males combine to drive the indigenous population to local extinction.

Mating interactions between closely related but reproductively isolated genetic groups are likely a common phenomenon (3, 22–24) and are expected given the widespread existence of hybridization and introgression (25). Although examples of asymmetric competition are well known (26–28), asymmetric mating interactions are less well described (28). The rarity of examples may be, as illustrated by this study, the consequence of such interactions leading to the rapid displacement of the disadvantaged organisms. Biological invasions offer opportunities to gauge and characterize the po-

tential magnitude and form of asymmetric mating interactions before species are lost through competitive exclusion or before the importance of competition is reduced over evolutionary time through niche partitioning and character displacement.

Allopatric species often demonstrate greater similarity in mating signals than do sympatric species, even when they have been diverging for a similar length of time (3). As a consequence of biological invasions, previously allopatric species are brought together and their partially similar mate recognition systems may promote asymmetric mating interactions between them. As we have shown, these interactions may play a critical role in determining the capacity of the invader to establish itself and the consequences for indigenous species.

References and Notes

1. J. L. Lookwood, M. F. Hoopes, M. P. Marchetti, *Invasion Ecology* (Blackwell, Malden, MA, 2007).
2. D. A. Holway, A. V. Suarez, *Trends Ecol. Evol.* **14**, 328 (1999).
3. R. Butlin, in *Speciation and the Recognition Concept, Theory and Application*, D. M. Lambert, H. G. Spencer, Eds. (Johns Hopkins Univ. Press, Baltimore, 1995), pp. 327.
4. J. K. Brown, D. R. Frohlich, R. C. Rosell, *Annu. Rev. Entomol.* **40**, 511 (1995).
5. L. M. Boykin *et al.*, *Mol. Phylogenet. Evol.* **44**, 1306 (2007).
6. M. Jiu *et al.*, *PLoS ONE* **2**, e182 (2007).
7. P. J. De Barro, J. W. H. Trueman, D. R. Frohlich, *Bull. Entomol. Res.* **95**, 193 (2005).
8. T. M. Perring *et al.*, *Science* **259**, 74 (1993).
9. International Union for the Conservation of Nature and Natural Resources (IUCN) list (www.issg.org).
10. R. Rekha, M. N. Maruthi, V. Muniyappa, J. Colvin, *Entomol. Exp. Appl.* **117**, 221 (2005).
11. T. M. Perring, in *Bemisia 1995: Taxonomy, Biology, Damage, Control and Management*, D. Gerling, R. T. Mayer, Eds. (Intercept, Andover, UK, 1996), p. 3–16.
12. L. H. C. Lima *et al.*, *Genet. Mol. Biol.* **25**, 217 (2002).
13. L. S. Zang, W. Q. Chen, S. S. Liu, *Entomol. Exp. Appl.* **121**, 221 (2006).

14. P. J. De Barro, P. J. Hart, *Bull. Entomol. Res.* **90**, 103 (2000).
15. P. J. De Barro, A. Bourne, S. A. Khan, V. A. L. Brancatini, *Biol. Invas.* **8**, 287 (2006).
16. C. Luo *et al.*, *Acta Entomol. Sin.* **45**, 759 (2002).
17. R. V. Gunning *et al.*, *J. Aust. Entomol. Soc.* **34**, 116 (1995).
18. See supporting material on Science Online.
19. L. S. Zang, S. S. Liu, *J. Insect Behav.* **20**, 157 (2007).
20. D. N. Byrne, T. S. Bellows, *Annu. Rev. Entomol.* **36**, 431 (1991).
21. Y. M. Ruan, J. B. Luan, L. S. Zang, S. S. Liu, *Entomol. Exp. Appl.* **124**, 229 (2007).
22. D. K. McInnis, D. J. Shure, *Oikos* **49**, 291 (1987).
23. B. Takafuji, E. Kuno, H. Fujimoto, *Exp. Appl. Acarol.* **21**, 379 (1997).
24. J. Yoshimura, W. T. Starmer, *Res. Popul. Ecol. (Kyoto)* **39**, 191 (1997).
25. J. M. Rhymer, D. Simberloff, *Annu. Rev. Ecol. Syst.* **27**, 83 (1996).
26. J. H. Lawton, M. P. Hassell, *Nature* **289**, 793 (1981).
27. R. F. Denno, M. S. McClure, J. R. Ott, *Annu. Rev. Entomol.* **40**, 297 (1995).
28. S. R. Reitz, J. T. Trumble, *Annu. Rev. Entomol.* **47**, 435 (2002).
29. We thank J. Liu, L. Zhang, K.-K. Lin, Z.-Y. Xiao, and G.-H. Yan for technical assistance, and J. Colvin, M. Dicke, A. Hoffmann, J. Trumble, T. Turlings, and M. Zalucki for comments and discussion. Supported by the National Basic Research and Development Program of China, National Natural Science Foundation of China, Ministry of Education of China, CSIRO Entomology, Horticulture Australia, and the Grains and Cotton Research and Development Corporations.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1149887/DC1

Materials and Methods

SOM Text

Figs. S1 to S4

Tables S1 to S5

References

29 August 2007; accepted 11 October 2007

Published online 8 November 2007;

10.1126/science.1149887

Include this information when citing this paper.

Declining Wild Salmon Populations in Relation to Parasites from Farm Salmon

Martin Krkošek,^{1,2†} Jennifer S. Ford,³ Alexandra Morton,⁴ Subhash Lele,¹ Ransom A. Myers,^{3*} Mark A. Lewis^{1,2}

Rather than benefiting wild fish, industrial aquaculture may contribute to declines in ocean fisheries and ecosystems. Farm salmon are commonly infected with salmon lice (*Lepeophtheirus salmonis*), which are native ectoparasitic copepods. We show that recurrent louse infestations of wild juvenile pink salmon (*Oncorhynchus gorbuscha*), all associated with salmon farms, have depressed wild pink salmon populations and placed them on a trajectory toward rapid local extinction. The louse-induced mortality of pink salmon is commonly over 80% and exceeds previous fishing mortality. If outbreaks continue, then local extinction is certain, and a 99% collapse in pink salmon population abundance is expected in four salmon generations. These results suggest that salmon farms can cause parasite outbreaks that erode the capacity of a coastal ecosystem to support wild salmon populations.

The decline in ocean fisheries (1, 2) and rise in global demand for fish have driven the rapid growth of aquaculture (3, 4). Although aquaculture may augment fish supply

(3), there are ecological risks, including competition and interbreeding of escaped farm fish with wild fish (5, 6), depletion of wild fish caught to feed farm fish (3, 4), and the spread of infec-

tion from farm fish to wild fish (7, 8). Disease threats of aquaculture to wild fish populations have long been contentious because of the uncertainty in impacts on those populations (9–12). We assess the impact of recurrent aquaculture-induced salmon lice (*L. salmonis*) infestations on wild pink salmon (*O. gorbuscha*) populations.

The salmon louse is a native marine ectoparasitic copepod of salmonids that feeds on surface tissues and causes stress, osmotic failure, viral or bacterial infection, and ultimately death (13). Lice are directly transmitted via planktonic nauplii and copepodids that can persist for several days. In areas without salmon farms, the prevalence of *L. salmonis* on juvenile pink salmon 2 to 3 months after marine emergence is low (<5%) (14–16), because returning adult salmon are mostly offshore when juvenile salmon enter the sea (16, 17). Louse infestations of wild juvenile salmon have occurred throughout the Broughton Archipelago in Pacific Canada (Fig. 1) from 2001 to 2005 (7, 8, 14, 18, 19). There, salmon farms situated in inlets and channels near rivers can increase copepodid densities above background levels for more than 80 km of wild salmon migration routes or, equivalently, for the first 2.5 months of the wild salmon's marine life (8). In response to a pink salmon population collapse in 2002, a primary migration corridor was fallowed in 2003 (i.e., farm salmon were removed from aquaculture facilities in Tribune Channel through Fife Sound, but farms peripheral to this route remained active) (Fig. 1). For that salmon cohort, *L. salmonis* abundance declined (19), and pink salmon marine survival increased (20).

To test for effects of lice on salmon population dynamics, we compiled Fisheries and Oceans Canada escapement data (the number of salmon per river), from 1970 to the present, for all pink salmon populations from rivers in the central coast of British Columbia, Canada (Fig. 1). There were 64 rivers whose salmon populations were not exposed to salmon farms and 7 rivers whose salmon populations must migrate past at least one salmon farm. Because pink salmon have a 2-year life cycle, there are distinct odd- and even-year lineages (21), which amount to 128 unexposed populations and 14 exposed populations. Rivers with substantial enhancement (e.g., spawning channels) were excluded because any increased salmon abundances in these rivers confound our estimates of natural changes in abundance. Unexposed populations had been and continue to be commercially fished. Exposed populations were commercially fished before the infestations, but the fishery

remains closed since the onset of the infestations, when the data show a marked decline in productivity (Fig. 2 and fig. S1).

The analysis was based on the Ricker model (22), which is commonly used to model time-series data from density-dependent populations (23–26), including pink salmon (24, 26), and provides robust estimates of population growth rates (24). The model is $n_i(t) = n_i(t-2)\exp[r - bn_i(t-2)]$, where $n_i(t)$ is the abundance of population i in year t , r is the population growth rate, and b determines density-dependent mortality. Upon log transformation to $\log[n_i(t)/n_i(t-2)] = r - bn_i(t-2)$, the Ricker equation becomes a linear model with intercept r and slope b that can be estimated by linear

regression and hierarchical mixed-effects modeling (23, 24, 27, 28). A preliminary model selection analysis did not support including random effects on r or b (fig. S2 and tables S1 and S2) (27). We therefore pooled data from multiple populations (27) and used linear regression to estimate parameters and parametric bootstrapping to construct 95% confidence intervals (CIs) on the parameter estimates (23). This allowed us to statistically compare parameters from pooled populations subjected to infestations, which is not possible with hierarchical mixed-effects models because there are only two data points per population during infestation years.

We compared parameter estimates among three groups: unexposed populations, exposed




Fig. 1. Study area in the Broughton Archipelago (boxed area in inset), depicting pink salmon populations from unexposed rivers (numbered circles) and exposed rivers (directly labeled within the lower rectangular frame). Inferred migration routes in the Broughton Archipelago are shown by the small arrows. Salmon farms are shown by black dots and sample sites by stars. Salmon farms south of Knight Inlet are not shown. Identities of the numbered (unexposed) rivers are provided in data set S1 (28).

¹Centre for Mathematical Biology, Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, AB, Canada. ²Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada. ³Biology Department, Dalhousie University, Halifax, NS, Canada. ⁴Salmon Coast Field Station, Simoom Sound, BC, Canada.

*Deceased.

†To whom correspondence should be addressed. E-mail: mkrkosek@ualberta.ca


preinfestation populations, and exposed populations during infestations (excluding the fallow year). The groups did not differ in b , and so we reanalyzed the data with b fixed among the three groups. Unexposed populations did not differ from exposed preinfestation populations in growth rate (unexposed populations: $r = 0.62$, 95% CI: 0.55 to 0.69; exposed preinfestation populations: $r = 0.68$, 95% CI: 0.46 to 0.90). The growth rate of exposed populations during the infestations was significantly lower and significantly negative ($r = -1.17$, 95% CI: -1.71 to -0.59 ; Fig. 3), meaning that if infestations are sustained, then local extinction is certain (29). Population viability analysis (28, 29) revealed the mean time to 99% population collapse is 3.9 generations, with the 95% CI from 3.7 to 4.2. During two generations of infestations, some exposed populations have declined to $<1\%$, whereas others have exceeded their historical abundance. We initially excluded the fallow data, because they contain only 1 year of observations and correspond to a nonrandom management action. By fixing $b = 0.64$, as estimated above, and estimating r from the remaining seven data points, we found the growth rate of fallow populations was significantly increased ($r = 2.50$, 95% CI: 1.28 to 3.62). The maximum reproductive rate for pink salmon is $r^* = 1.2$ (24). Fishing mortality probably reduced r for unexposed and exposed preinfestation populations. The depressed growth rate of exposed salmon populations during the infestations indicates that previous fishing mortality (now ceased) has been greatly exceeded by louse-induced mortality.

To estimate the mortality of pink salmon caused by lice, we extended the Ricker model to directly accommodate louse data collected from exposed populations during the infestations (14, 18, 19, 28). We constrained the model by fixing $b = 0.64$ and by requiring $r = r^* = 1.2$, because there was no fishing mortality. Louse-induced mortality is represented by multiplying by $\exp[-aP_i(t-1)]$, where P is the mean abundance of motile (adult and preadult) lice per juvenile salmon from population i that spawned

in year t . We log-transformed the model to $\log[n_i(t)/n_i(t-2)] = r - bn_i(t-2) - aP_i(t)$ and used linear regression to estimate a . The term $\exp[-aP_i(t-1)]$ significantly improved the fit of the model ($t = -5.019$, $df = 33$, $P = 1.74 \times 10^{-5}$; fig. S3), and results remained strong when the data were restricted by averaging populations and excluding some population groups ($P < 0.005$ for all groups; table S3). The parameter a corresponds to the rate of parasite-induced host mortality multiplied by the time that juvenile salmon are exposed to the parasites, $a = \alpha T$. The exposure time, T , is about 2 months (based on the migration speed of juvenile pink salmon through the archipelago), and the value of α has been estimated at 0.022 (motile lice \times day) $^{-1}$ (based on survival experiments of naturally infected juvenile pink salmon) (8). Dividing the estimated $a = 0.89$ (95% credible intervals are from 0.46 to 1.34) by 60 days reveals an excellent correspondence between these two independent estimates of pathogenicity ($a/60 = 0.015$, with 95% credible estimates from 0.0077 to 0.022). Using a hierarchical Bayesian simulation (28) that represents uncertainty in the model fit as well as in the distribution of r^* (12), we found the estimated mortality of pink salmon, $1 - \exp[-aP_i(t-1)]$, caused by lice ranged from 16% to over 97% and was commonly over 80% (Table 1). The lowest mortality comes from fallow populations when louse abundance was nevertheless elevated, possibly resulting from transmission from active farms outside the fallowed corridor (7, 19, 20).

These results provide strong empirical evidence that salmon farm-induced *L. salmonis* infestations of juvenile pink salmon have depressed wild pink salmon populations and may lead to their local extinction. However, this parasite threat may not exist at low farm salmon abundances; the delay between the onset of salmon aquaculture in 1987 and louse infestations in 2001 (Fig. 2) may be explained by farm fish abundance crossing a host density threshold above which outbreak conditions occur (30). It is unlikely that another factor is responsible: The

Fig. 2. Time series of normalized population deviances $\{\log[N_i(t)/m_i]\}$, where $N_i(t)$ is the population estimate for population i in year t and m_i is the time-series mean abundance for population i for 128 control populations of pink salmon (open gray circles) and 14 pink salmon populations exposed to salmon farms (black circles). The vertical dashed line marks the beginning of salmon aquaculture in the Broughton Archipelago. The vertical solid line marks the onset of louse infestations (and the commercial fishery closure) affecting the exposed populations. The arrow indicates data for exposed pink salmon cohorts that, as juveniles, experienced a fallowed migration corridor.



increased growth rate in response to following rules out other factors that could have affected exposed, but not unexposed, populations. The results rely on extensive spatial replication to compensate for short time series in infestation




Fig. 3. Fits of the log-transformed Ricker model to escapement data for unexposed populations (A), exposed populations before infestations (B), and exposed populations during the infestations (C), and a comparison of the log-transformed Ricker model for the three groups in panels (A) to (C) (D). The intercept (growth rate) is lower for the exposed population during the infestations than for exposed populations before the infestations and the unexposed populations.

Table 1. Mean abundances, P , of motile *L. salmonis* on juvenile pink salmon and estimated parasite-induced host mortality, M (with upper and lower bounds of the 95% credible interval in parentheses), for exposed populations during infestations.

River	2002		2003		2004*		2005		2006	
	P	M	P	M	P	M	P	M	P	M
Ahta	3.4	95.21 (79.07, 98.95)	1.0	59.09 (36.87, 73.82)	0.3	23.52 (12.89, 33.10)	2.6	90.21 (69.76, 96.93)	0.4	30.06 (16.81, 41.49)
Kakweiken	3.4	95.21 (79.07, 98.95)	1.0	59.09 (36.87, 73.82)	0.3	23.52 (12.89, 33.10)	2.6	90.21 (69.76, 96.93)	0.4	30.06 (16.81, 41.49)
Viner	4.0	97.20 (84.12, 99.53)	2.2	86.00 (63.65, 94.76)	0.2	16.37 (8.79, 23.51)	2.3	87.20 (65.29, 95.41)	1.4	71.39 (47.48, 84.68)
Wakeman	4.0	97.20 (84.12, 99.53)	2.2	86.00 (63.65, 94.76)	0.2	16.37 (8.79, 23.51)	2.3	87.20 (65.29, 95.41)	1.4	71.39 (47.48, 84.68)
Kingcome	4.0	97.20 (84.12, 99.53)	2.2	86.00 (63.65, 94.76)	0.2	16.37 (8.79, 23.51)	2.3	87.20 (65.29, 95.41)	1.4	71.39 (47.48, 84.68)
Ahnuhati	2.6	90.21 (69.76, 96.93)	0.7	46.51 (27.53, 60.86)	0.2	16.37 (8.79, 23.51)	1.9	81.70 (58.27, 92.16)	0.3	23.52 (12.89, 33.10)
Lull	2.6	90.21 (69.76, 96.93)	0.7	46.51 (27.53, 60.86)	0.2	16.37 (8.79, 23.51)	1.9	81.70 (58.27, 92.16)	0.3	23.52 (12.89, 33.10)

*These data correspond to the salmon cohort responding to the fallow treatment in 2003.

years. The time to reach sufficient temporal replication to support hierarchical mixed-effects modeling, say 10 generations (which equals 20 years), greatly exceeds the predicted time to extinction. That is, there is a major risk associated with waiting for large data sets to accumulate before implementing conservation policy. Industrial aquaculture is rapidly expanding to new species, regions, and habitats (31), which can create parasite outbreaks that contribute to the decline of ocean fisheries and ecosystems.

References and Notes

- J. B. C. Jackson *et al.*, *Science* **293**, 629 (2001).
- R. A. Myers, B. Worm, *Nature* **423**, 280 (2003).
- R. L. Naylor *et al.*, *Nature* **405**, 1017 (2000).
- R. Goldberg, R. Naylor, *Front. Ecol. Environ.* **3**, 21 (2005).
- I. A. Fleming *et al.*, *Proc. R. Soc. London Ser. B* **267**, 1517 (2000).
- J. P. Volpe, B. R. Anholt, B. W. Glickman, *Can. J. Fish. Aquat. Sci.* **58**, 197 (2001).
- M. Krkošek, M. A. Lewis, J. P. Volpe, *Proc. R. Soc. London Ser. B* **272**, 689 (2005).
- M. Krkošek, M. A. Lewis, A. Morton, L. N. Frazer, J. P. Volpe, *Proc. Natl. Acad. Sci. U.S.A.* **103**, 15506 (2006).
- A. H. McVicar, *ICES J. Mar. Sci.* **54**, 1093 (1997).
- D. J. Noakes, R. J. Beamish, M. L. Kent, *Aquaculture* **183**, 363 (2000).
- A. H. McVicar, *Aquac. Res.* **35**, 751 (2004).
- R. Hilborn, *Proc. Natl. Acad. Sci. U.S.A.* **103**, 15277 (2006).
- A. W. Pike, S. L. Wadsworth, *Adv. Parasitol.* **44**, 233 (2000).
- A. Morton, R. Routledge, C. Peet, A. Ladwig, *Can. J. Fish. Aquat. Sci.* **61**, 147 (2004).
- C. R. Peet, thesis, University of Victoria (2007).
- M. Krkošek *et al.*, *Proc. R. Soc. London Ser. B* **274**, 3141 (2007).
- C. Groot, L. Margolis, Eds. *Pacific Salmon Life Histories* (UBC Press, Vancouver, 1991).
- A. B. Morton, R. Williams, *Can. Field Nat.* **117**, 634 (2003).
- A. Morton, R. D. Routledge, R. Williams, *N. Am. J. Fish. Manage.* **25**, 811 (2005).
- R. J. Beamish *et al.*, *ICES J. Mar. Sci.* **63**, 1326 (2006).
- W. R. Heard, in (17), pp. 119–230.
- W. E. Ricker, *J. Fish. Res. Board Can.* **11**, 559 (1954).
- B. Dennis, M. L. Taper, *Ecol. Monogr.* **64**, 205 (1994).
- R. A. Myers, K. G. Bowen, N. J. Barrowman, *Can. J. Fish. Aquat. Sci.* **56**, 2404 (1999).
- B. W. Brook, C. J. A. Bradshaw, *Ecology* **87**, 1445 (2006).
- F. J. Mueter, R. M. Peterman, B. J. Pyper, *Can. J. Fish. Aquat. Sci.* **59**, 456 (2002).
- J. C. Pinheiro, D. M. Bates, *Mixed-Effects Models in S and S-PLUS* (Springer, New York, 2004).
- Materials and methods are available as supporting material on Science Online.

- B. Dennis, P. L. Munholland, J. M. Scott, *Ecol. Monogr.* **61**, 115 (1991).
- J. O. Lloyd-Smith *et al.*, *Trends Ecol. Evol.* **20**, 511 (2005).
- C. M. Duarte, N. Marbá, M. Holmer, *Science* **316**, 382 (2007).
- We dedicate this paper to our coauthor, Ransom Myers, who passed away before the completion of this work. We thank J. Volpe, J. Reynolds, L. Dill, M. Wonham, B. Connors, and A. Gottesfeld for helpful comments; A. Park for assistance in preparing data and figures; and Fisheries and Oceans Canada stock assessment scientists who collected and shared data. Funding came from the Natural Science and Engineering Research Council of Canada, the Canadian Mathematics of Information Technology and Complex Systems National Centre of Excellence Network on Biological Invasions and Dispersal

Research (with nonacademic participants including the David Suzuki Foundation, Canadian Sablefish Association, Wilderness Tourism Association, and Finest at Sea), the National Geographic Society, Tides Canada, a University of Alberta Bill Shostak Wildlife Award, the Lenfest Ocean Program, Census of Marine Life, and a Canada Research Chair.

Supporting Online Material

www.sciencemag.org/cgi/content/full/318/58571772/DC1
Materials and Methods
Figs. S1 to S3
Tables S1 to S3
Dataset S1

2 August 2007; accepted 2 November 2007
10.1126/science.1148744

Habitat Split and the Global Decline of Amphibians

Carlos Guilherme Becker,^{1,2} Carlos Roberto Fonseca,^{2*} Célio Fernando Baptista Haddad,³ Rômulo Fernandes Batista,⁴ Paulo Inácio Prado⁵

The worldwide decline in amphibians has been attributed to several causes, especially habitat loss and disease. We identified a further factor, namely “habitat split”—defined as human-induced disconnection between habitats used by different life history stages of a species—which forces forest-associated amphibians with aquatic larvae to make risky breeding migrations between suitable aquatic and terrestrial habitats. In the Brazilian Atlantic Forest, we found that habitat split negatively affects the richness of species with aquatic larvae but not the richness of species with terrestrial development (the latter can complete their life cycle inside forest remnants). This mechanism helps to explain why species with aquatic larvae have the highest incidence of population decline. These findings reinforce the need for the conservation and restoration of riparian vegetation.

Amphibian populations are declining worldwide (1, 2). Among the factors determining the amphibian declines are habitat loss and fragmentation, which affect amphibians just as they affect any other organisms: through population isolation, inbreeding, and edge effects (3–5). Another important factor is the fungus *Batrachochytrium dendrobatidis*, a highly virulent pathogen that attacks many amphibian species and has been responsible for the decline of many populations even in undisturbed environments (6, 7). Amphibians can also be threatened by climate shifts (7), ultraviolet-B radiation (8), introduction of exotic species (9), and agrochemical contaminants (10). We inves-

tigated the role of a further factor, which we define as “habitat split.”

Amphibian species with aquatic larvae typically undergo a major ontogenetic niche shift, whereby tadpoles and adults occupy two distinct habitats (11). In pristine environments, the aquatic habitat of the tadpoles and the terrestrial habitat of the postmetamorphics grade into each other. However, in landscapes occupied by humans, land use has often resulted in a spatial separation between remnants of terrestrial habitat and breeding sites (12). Adults of species with aquatic larvae, in order to breed, are obliged to abandon forest remnants to reach water bodies, and at the end of the reproductive season, both

ERRATUM

Post date 19 December 2008

Reports: “Declining wild salmon populations in relation to parasites from farm salmon” by M. Krkošek *et al.* (14 December 2007, p. 1772). This correction summarizes small changes to the statistical results written in the main text of the Report on the effects of sea lice infestations on pink salmon population dynamics. Small changes have also been made to data set S1 and tables S2 and S3 in the Supporting Online Material. The changes to the statistical results do not affect the conclusions of the report.

The changes arise due to revisions of 11 escapement estimates for exposed populations that were not present in the original data provided to the authors by the Canadian Department of Fisheries and Oceans. The changes have been confirmed by Brian Riddell, Division Head, Salmon Assessment and Freshwater Ecosystems, Pacific Biological Station, Fisheries and Oceans Canada.

Table 1.

Point estimates and 95% confidence intervals for the population growth rate r from the Ricker model fit to grouped pink salmon escapement data.

Data set used	Group	r	95% confidence interval (CI)
Original	Unexposed	0.62	0.55 to 0.69
	Exposed pre-lice	0.68	0.46 to 0.90
	Exposed infested	-1.17	-1.71 to -0.59
	Exposed fallow	2.50*	1.28 to 3.62*
Corrected	Unexposed	0.63	0.56 to 0.70
	Exposed pre-lice	0.70	0.47 to 0.92
	Exposed infested	-1.16	-1.68 to -0.62
	Exposed fallow	2.63	1.39 to 3.78

*Results shown as reported in the paper. This should actually read $r = 2.63$ with 95% CI of 1.38 to 3.77.

Population growth rates. The population growth rate r was estimated from the Ricker model for four groups of data. There are small changes to the point estimates of r as well as the 95% confidence intervals. The changes are summarized in Table 1. The associated estimate of b for density-dependent mortality has changed from its original value of 0.64 to its corrected value 0.65.

Population viability analysis. In this section, a population viability analysis was applied to pink salmon populations in the Broughton Archipelago during sea lice infestation years. Small changes to the results are summarized in Table 2.

Continued on next page

ERRATUM

Post date 19 December 2008

Louse-induced salmon mortality. The Ricker model was extended to test whether including louse-induced mortality of wild pink salmon improved the fit of the model. The analysis consisted of estimating a parameter a . The point estimate for the parameter has changed from 0.89 to 0.90. The 95% credible intervals for the parameter a from the analysis using the unconstrained data changed from 0.46 to 1.34 in the original analysis to 0.47 to 1.34 using the corrected data set.

Table 2.

Population viability analysis of Broughton Archipelago pink salmon populations subjected to sea lice infestations.		
Data set used		
Original	Population growth rate	-1.17
	Variance of environmental stochasticity	1.92
	Mean time to 99% collapse	3.9 (95% CI 3.7 to 4.2)
Corrected	Population growth rate	-1.16
	Variance of environmental stochasticity	1.90
	Mean time to 99% collapse	4.0 (95% CI 3.7 to 4.2)